


REACTOR STUDIES FOR THE LHD AND THE MHH2


P.R. Garabedian

The NSTAB code has been run to show that the LHD stellarator is linearly unstable, but remains nonlinearly stable, at the β of 3.2% achieved experimentally. At observed temperatures of 3 keV the TRAN code predicts an energy confinement time of 160 ms that agrees with measured values. Predictions of ballooning stability for the LHD are more pessimistic than estimates from bifurcated solutions calculated over 1, 2, 5 or 10 periods.


Good correlation of computations with observations in the LHD have been applied to assess results about equilibrium, stability, and transport for a quasiaxially symmetric MHH2 reactor. Twelve only moderately twisted coils provide robust magnetic surfaces. The β limit is 4%, and 12 only moderately twisted coils provide robust magnetic surfaces.




Poincaré map of the flux surfaces at four cross sections over the full torus of a bifurcated LHD equilibrium at $\beta = 0.025$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a standard pressure profile $p = p_0(1-s)$, and with bootstrap current, the global $m = 1, n = 1$ mode of this solution is linearly unstable, but nonlinearly stable.




Poincaré map of the flux surfaces at four cross sections over five field periods of a bifurcated LHD equilibrium at $\beta = 0.02$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a triangular pressure profile $p = p_0(1 - s^{0.7})$ like observed values of the electron temperature, the solution is linearly unstable but nonlinearly stable.




Poincaré map of the flux surfaces at four cross sections over five field periods of a bifurcated LHD equilibrium at $\beta = 0.03$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a triangular pressure profile $p = p_0(1 - s^{0.8})$ like observed values of the electron temperature, the solution is linearly unstable but nonlinearly stable.



Poincaré map of the flux surfaces at four cross sections over two field periods of a bifurcated LHD equilibrium at $\beta = 0.04$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a standard pressure profile $p = p_0(1 - s)$, this solution developed a ballooning mode and is becoming nonlinearly unstable. The mode was stable at $\beta = 0.03$.





Poincaré map of the flux surfaces at four cross sections over one field period of a bifurcated LHD equilibrium at $\beta = 0.04$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a standard pressure profile $p = p_0(1 - s)$, this solution developed a ballooning mode and is becoming nonlinearly unstable. The mode was stable at $\beta = 0.03$.




NSTAB calculation of the flux surfaces at four cross sections over one field period for an LHD equilibrium at $\beta = 0.02$ with the magnetic axis shifted inward to a position with plasma radius $R = 3.6$. For a standard pressure profile $p = p_0(1 - s)$, this solution remains both linearly and nonlinearly stable after perturbation by low order ballooning modes.

L = poloidal transit







Four cross sections of the flux surfaces over one field period of a bifurcated W7-AS equilibrium at $\beta = 0.03$ with a pressure profile $p = p_0(1 - s)$ and with net current bringing the rotational transform into the interval $0.6 > \iota > 0.5$. At conditions related to those in the experiment an ideal MHD mode with ballooning structure appears in the calculation. After the net current was removed the solution became stable. In this example the NSTAB code was run 10^5 cycles using spectral terms of degrees 32 and 24 in the poloidal and toroidal angles




Cycles of a calculation of the energy confinement time τ_E in milliseconds for an NBI shot of the LHD experiment using a quasineutrality algorithm to adjust the electric potential Φ . Oscillations of Φ along the magnetic lines model turbulence and anomalous transport, so there is good agreement with the measured value.




Iterations to quasineutrality in a Monte Carlo study of the energy confinement time τ_E , measured in milliseconds, for an LHD stellarator with major radius 25 m and plasma radius 4.5 m at reactor conditions with average $T = 12 \text{ keV}$, $n = 1.2 \times 10^{14} \text{ cm}^{-3}$, and $B = 5 \text{ tesla}$. Transport is good because the radial electric field has risen to a potential level four times as big as the temperature.




Iterations to quasineutrality in a Monte Carlo computation of the energy confinement time τ_E , measured in milliseconds, for an MHH2 stellarator with major radius 10 m and plasma radius 3 m at reactor conditions with average $T = 12$ keV, $n = 1.7 \times 10^{14} \text{ cm}^{-3}$, and $B = 5$ tesla. The magnetic spectrum has good axial symmetry, but we have superimposed a mirror term anyway that improves transport by driving the radial electric field to a potential level four times as big as the temperature.




Four cross sections of the flux surfaces over the full torus of a stable MHH2 equilibrium at $\beta = 0.04$ with a pressure profile $p = p_0(1 - s^{1.5})^{1.5}$ and with net current bringing the rotational transform into the interval $0.48 > \iota > 0.45$. In this example the NSTAB code was run 35,000 cycles using spectral terms of degree 24 in each of the poloidal and toroidal angles and using 27 radial mesh points.




Four Poincaré sections of the flux surfaces over two field periods of an MHH2 equilibrium with a pressure profile $p = p_0(1 - s^2)^2$, a β of 5%, and $0.6 \geq \iota \geq 0.4$. During a run of 5×10^5 cycles of the NSTAB code using trigonometric polynomials of degree 20, an asymmetrical ballooning mode converged to a symmetric pair of bulges in regions of bad curvature of a wall stabilized solution of the magnetostatic equations that minimizes E .



Six out of twelve modular coils of the MHH2 stellarator in a vacuum magnetic field given by the Biot-Savart law. Judicious filtering of the Fourier series used to calculate filaments specifying the geometry of the configuration defines shapes that are not excessively twisted. Parameters have been adjusted to provide ample space around each coil, and the aspect ratio of the plasma is 3.5.



Poincaré section of magnetic surfaces for the MHH2 stellarator displaying the control surface for the coils, the known shape of the plasma at $\beta = 0$, and magnetic lines computed in a plasma with five big islands where the rotational transform crosses the resonant value $\iota = 2/5$.



Poincaré section of magnetic surfaces for the MHH2 stellarator displaying the control surface for the coils, the known shape of the plasma at $\beta = 0$, and magnetic lines computed in a plasma with islands that have been suppressed where the rotational transform crosses the resonant value $\iota = 2/5$.

$$\int\int\int(\frac{1}{2}B^2-p)\,dV\,=\,\text{minimum}$$


$$\mathbf{B}=\nabla s\times\nabla\theta=\nabla\phi-\zeta\nabla s\,\,,\,\,\mathbf{J}=\nabla s\times\nabla\zeta$$

$$r+iz=e^{iu}\sum\Delta_{mn}\,e^{-imu+inv}$$

$$\frac{1}{B^2}=\sum B_{mn}\cos(m\theta-[n-\imath m]\phi)$$

$$\zeta=p'\sum\frac{B_{mn}}{n-\imath m}\sin(m\theta-[n-\imath m]\phi)$$

$$\mathbf{B}=\nabla\times\sum_{j=0}^k\delta s_j\int\int\frac{\nabla\zeta_j\times\mathbf{N}}{4\pi r}d\sigma$$



Poincaré section of a free boundary calculation for the MHH2 stellarator displaying the control surface for the coils, the known shape of the plasma at $\beta = 0$, filaments that simulate currents in the plasma driven by the pressure, and magnetic lines computed at finite β in the scrape-off layer of the solution, where the rotational transform is crossing the resonant value $\iota = 2/5$.